



Parachorius semsanganus sp. n. (Coleoptera, Scarabaeidae, Scarabaeinae) from Laos and its significance in the phylogeny of Oriental Deltotchilini

Tarasov, Sergey; Keith, D.

Published in:
ZooKeys

DOI:
[10.3897/zookeys.111.1221](https://doi.org/10.3897/zookeys.111.1221)

Publication date:
2011

Document version
Publisher's PDF, also known as Version of record

Document license:
[CC BY](#)

Citation for published version (APA):
Tarasov, S., & Keith, D. (2011). *Parachorius semsanganus* sp. n. (Coleoptera, Scarabaeidae, Scarabaeinae) from Laos and its significance in the phylogeny of Oriental Deltotchilini. *ZooKeys*, 111, 51-57.
<https://doi.org/10.3897/zookeys.111.1221>

***Parachorius semsanganus* sp. n. (Coleoptera, Scarabaeidae, Scarabaeinae) from Laos and its significance in the phylogeny of Oriental Deltochilini**

S. Tarasov^{1,2,†}, D. Keith^{3,‡}

1 Department of Entomology, Natural History Museum of Denmark/University of Copenhagen, Zoological Museum, Universitetsparken 15, DK-2100 Copenhagen, Denmark **2** Institute of Natural Science, Kaluga State University, Stepana Razina str. 26, Kaluga 248023, Russia **3** Muséum des Sciences Naturelles et de Préhistoire, 5 bis, boulevard de la Courtille, 28000 Chartres, France

† [urn:lsid:zoobank.org:author:E8121729-4706-476B-BA34-64CC2CA02855](https://zoobank.org/urn:lsid:zoobank.org:author:E8121729-4706-476B-BA34-64CC2CA02855)

‡ [urn:lsid:zoobank.org:author:C922B63D-331E-44BF-B386-9A70AE05CA28](https://zoobank.org/urn:lsid:zoobank.org:author:C922B63D-331E-44BF-B386-9A70AE05CA28)

Corresponding author: S. Tarasov (sergxf@yandex.ru)

Academic editor: Andrey Frolov | Received 10 March 2011 | Accepted 5 May 2011 | Published 22 June 2011

[urn:lsid:zoobank.org:pub:45135830-3DBF-4567-9324-9E6B06174B93](https://zoobank.org/pub:45135830-3DBF-4567-9324-9E6B06174B93)

Citation: Tarasov S, Keith D (2011) *Parachorius semsanganus* sp. n. (Coleoptera, Scarabaeidae, Scarabaeinae) from Laos and its significance in the phylogeny of Oriental Deltochilini. ZooKeys 111: 51–57. doi: 10.3897/zookeys.111.1221

Abstract

The new species *Parachorius semsanganus* sp. n. is described from Laos. This enigmatic Oriental deltochiline represents a “morphological link” between *Parachorius* and *Cassolus* by sharing characters of the two genera. The fact that *P. semsanganus* cannot be unequivocally placed in either of these two genera stresses some more general problems of the current classification of *Parachorius* and *Cassolus*. Such problems can be solved only in the course of phylogenetic analysis, the need of which is briefly outlined.

Keywords

Deltochilini, *Parachorius*, *Cassolus*, new species, Laos, “morphological link”

Introduction

The dung beetle tribe Deltotrichini Lacordaire, 1856 (= Canthonini Lansberge, 1874, synonymy according to Bouchard et al. (2011)), displays a Gondwanian distribution and is the most genus-rich tribe of the Scarabaeinae. It comprises 873 species grouped into 91 genera that constitute 40% of the entire generic diversity of the subfamily Scarabaeinae (Davis et al. 2008). However, the monophyly of the tribe is doubtful according to a revision of African genera (Scholtz and Howden 1987), and it appears polyphyletic in the morphology-based and molecular phylogenies (Philips et al. 2004; Ocampo and Hawks 2006; Monaghan et al. 2007; Sole and Scholtz 2010). Such ambiguity highly complicates the diagnosis of the tribe and thus makes a consensus about its generic composition impossible at the moment.

The Oriental genus *Parachorius* Harold, 1873 comprises six species and, according to the last monographic study (Balthasar 1963), belongs to the tribe Pinotini Kolbe, 1905, which is a synonym of the Ateuchini Laporte, 1840 (Smith 2006; Bouchard et al. 2011). Vaz-de-Mello (2007; 2008) places *Parachorius* within the tribe Coprini Leach, 1815 based on morphological phylogenetic analysis of Ateuchini and related taxa. By contrast, another preliminary morphology-based phylogeny of Oriental Deltotrichini using aedeagal and somatic characters (Tarasov, unpublished) strongly suggests that *Canthon* Hoffmannsegg, 1817 (the most speciose genus of the Deltotrichini), *Parachorius*, and another Oriental deltochiline genus *Cassolus* Sharp, 1875, form a monophyletic group. Therefore, here, we treat *Parachorius* as a deltochiline. Based on study of external morphological characters within the tribe Deltotrichini, *Parachorius* seems to be most closely related to the genus *Cassolus*. The genus *Cassolus* includes nine species that are morphologically very heterogeneous. Such morphological variation, especially when compared with the morphology of the known species of *Parachorius*, may indicate a nested position of the latter genus within *Cassolus* (Tarasov, unpublished).

Whilst surveying recent scarab collections from Laos deposited at the Naturhistorisches Museum in Basel, Switzerland (NHMB), we discovered a very interesting deltochiline species displaying a mixed character set between *Parachorius* and *Cassolus*. This species cannot be unequivocally placed in either genus using current taxonomic concepts of these taxa (Balthasar 1963). However, we tentatively place this new species in the genus *Parachorius*.

Discovery of this species supports the above mentioned evidence for the close relationship between *Cassolus* and *Parachorius* as well as our provisory placement of *Parachorius* within the Deltotrichini. Its description, provided here, enables its incorporation in the upcoming phylogenetic analysis of the entire generic complex.

Material and methods

All photos were taken with a digital camera attached to a dissecting microscope (Leica MZ16A). Male aedeagi in Figs 1–3 were photographed in glycerin. First, the dissected

aedeagus was macerated in 10% solution of KOH for several hours and then rinsed with distilled water. Finally, the aedeagus was placed in glycerin for taking pictures and subsequent storage.

All the material used in this study is housed in the NHMB.

Species description

Parachorius semsanganus Tarasov & Keith *sp. n.*

urn:lsid:zoobank.org:act:E79FA444-725E-40E6-986A-9FB48B7D232A

http://species-id.net/wiki/Parachorius_semsanganus

Figs 1–5

Type locality. Laos, Xieng Khouang prov., Phou Sane Mt.

Material examined. *Holotype* (NHMB), male bearing the following labels:

LAOS-NE, Xieng Khouang prov., 19°38.20'N 103°20.20'E, Phonsavan (30 km NE): PHOU SANE Mt., 1420 m, 10.-30.v.2009, D. Hauck leg.

NHMB Basel, NMPC Prague Laos 2009 Expedition: M. Brancucci, M. Geiser, Z. Kraus, D. Hauck, V. Kubáň

HOLOTYPE *Parachorius semsanganus* S. Tarasov & D. Keith det. 2011

Paratypes. 9♀, same data as holotype; 2♂, same data as holotype but Z. Kraus leg.; 4♂, 3♀, LAOS-NE, Xieng Khouang prov., 19°37-8.'N 103°20'E, 30 km NE Phonsavan: Ban Na, Lam Phou Sane Mt., 1300–1500 m, 10.-30.v.2009, M. Brancucci leg.


Description. Oval, convex, black, entirely shiny; mouthparts, antennae and legs red-brown. Dorsal body side covered with two types of punctures: larger (normal) punctures and very tiny punctures (which can be observed only under higher magnification of 40x or more) Dorsal and ventral body sides glabrous. Length 8.2–10.6

Male (Fig. 4). Head flat, punctuation fine; anterior margin notched medially; notch delimited by 2 prominent triangular obtuse teeth; clypeus laterad of each tooth very slightly notched; eyes completely divided by canthus into lower and upper lobes; lower lobes significantly larger than upper ones; genae and clypeus not distinctly separated from frons; genae rounded and protruding; antennae with 9 segments, antennal club with 3 segments.

Pronotum broadly trapezoidal, punctuation fine, separated by 1–2 puncture diameters on disc, becoming slightly denser laterally. Lateral margins of pronotum flattened, arcuate, widest near base; lateral and anterior side marginate, posterior side not marginate; anterior angles obtuse; posterior angles rounded. Prothoracic fovea excavated, delimited by ridge reaching propleural lateral margin.

Elytra with eight striae, sublateral carina forming pseudopipleuron beyond eighth stria; epipleura narrow; interstriae flat with sparse, fine punctuation.

Protibiae with three outer teeth; 1st tooth slightly thicker than two others; inner margin with two vertical teeth underneath, located approximately opposite to 2nd and



Figures 1–5. Morphological features of *Parachorius semsanganus* sp. n.: 1–3 paratype, aedeagus 1 aedeagus lateral view 2 aedeagus apical view 3 aedeagus dorsal view 4 male holotype, habitus 5 male paratype, hind leg, arrow indicates teeth on inner tibial margin.

3rd outer teeth; protibial apical spur acute, long, reaching middle or apical portion of 3rd tarsal segment; sometimes protibial teeth and apical spur abraded.

Metafemoral posterior margin with keel bearing indistinct and slight serration on top (Fig. 5). Metatibiae slightly curved, conspicuously denticulate on inner margin (Fig. 5, indicated with arrow); teeth are abraded in some specimens.

Pygidium with rather coarse, uniform, dense punctation.

Aedeagus (Figs 1–3) with converging, spatulate apices of parameres.

Female. Similar to male but with the 1st protibial outer tooth slightly thinner than in males; metafemoral posterior margin not serrate; metatibial inner margin not denticulate.

Variation. All specimens of the type series look very similar to each other. Some variation may be observed in the shape of teeth on the metatibial inner margin, which are less expressed in some males due to abrasion.

Holotype (Fig. 4). The holotype specimen lacks the tarsus of the right middle leg.

Differential diagnosis. The new species is quite distinct among all other known species of *Parachorius* and *Cassolus*. It can be easily separated from them by the following unique set of character states: clypeus near outer side of each clypeal tooth very slightly notched, metatibial inner margin with large teeth (Fig. 5, arrowed), and aedeagus with spatulate apices which are largely bent inward (Figs 1–3).

Distribution and ecology. The species is known from only 16 specimens of the type series collected across a range of altitudes between 1300–1500 m on Phou Sane Mt. of Xieng Khouang province in Laos.

Etymology. The name of the new species is derived from the Latinized Lao words “syan” – link and “sanga” – spectacular. Its meaning “spectacular link” refers to the fact that this species represents a “morphological link” between the genera *Parachorius* and *Cassolus*.

Taxonomic notes. Based on taxonomic concepts of the most recent monographic study dealing with *Parachorius* and *Cassolus* (Balthasar 1963), the morphological differences between these two taxa can be summarized as follows: *Cassolus* are normally smaller than *Parachorius*; the clypeus near the outer side of each clypeal tooth is usually deeply notched in *Cassolus* and not notched in *Parachorius* (very slightly notched in *P. semsanganus* sp. n.); the metatibiae are curved in *Cassolus* and more or less straight in *Parachorius* (slightly curved in *P. semsanganus* sp. n.); some *Cassolus* species have denticles on the inner metatibial margin, whereas the metatibial margin of *Parachorius* is not at all denticulate (distinctly denticulate in *P. semsanganus* sp. n.). As can be seen from this combination of characters, *P. semsanganus* sp. n. is similar, on the one hand, to *Parachorius* and on the other hand to *Cassolus* (in particular to *C. gotoi* Masumoto, 1986). A robust justification of the taxonomic placement of *P. semsanganus* sp. n. requires an extensive phylogenetic analysis embracing both *Parachorius* and *Cassolus*. Such an analysis is currently in preparation and it may, in particular, result in the synonymy of *Parachorius* and *Cassolus*. Therefore to avoid potential nomenclatural changes in the future, we place the new species in the earlier described genus *Parachorius*.

Acknowledgements

We are very thankful to Alexey Solodovnikov (Natural History Museum of Denmark, Copenhagen) for his suggestions, advice and first review of the manuscript. Adrian Davis (University of Pretoria, Republic of South Africa), Jan Krikken (Nationaal Natuurhistorisch Museum, Leiden, the Netherlands), Andrey Frolov (Zoological Institute, St. Petersburg, Russia) and Daniel Whitmore (Corpo Forestale dello Stato - Centro Nazionale Biodiversità Forestale “Bosco Fontana”, Verona, Italy) are sincerely acknowledged for their suggestions and corrections that significantly helped to improve this paper. Our thanks are due to Michel Brancucci and Isabelle Zürcher (NHMB) for making material under their care available for this study. We are also grateful to Pierre Tautzin (Vanves, France) for his help in obtaining needed literature, and to Tristao Branco (Porto, Portugal) for his advice and help. This paper is completed during the stay of one of us (S.T.) as a visiting student at the Department of Entomology of the Natural History Museum of Denmark, under financial support from the grants of that department. S.T. is sincerely thankful to friendly team of the Department for their help and support.

References

- Balthasar V (1963) Monographie der Scarabaeidae und Aphodiidae der palaearktischen und orientalischen Region. Tschechoslowakische Akademie der Wissenschaften Prag 1: 1–391, XXIV pl.
- Bouchard P, Bousquet Y, Davies A, Alonso-Zarazaga M, Lawrence J, Lyal C, Newton A, Reid C, Schmitt M, Slipinski A, Smith A (2011) Family-Group Names In Coleoptera (Insecta). *Zookeys* 88: 1–972. doi:10.3897/zookeys.88.807
- Davis A, Frolov A, Scholtz C (2008) The African Dung Beetle genera. Protea Book House, 272 pp.
- Monaghan MT, Inward DJG, Hunt T, Vogler AP (2007) A molecular phylogenetic analysis of the Scarabaeinae (dung beetles). *Molecular Phylogenetics and Evolution* 45: 674–692. doi:10.1016/j.ympev.2007.06.009
- Ocampo FC, Hawks DC (2006) Molecular phylogenetics and evolution of the food relocation behaviour of the dung beetle tribe Eucraniini (Coleoptera: Scarabaeidae: Scarabaeinae). *Invertebrate Systematics* 20: 557–570. doi:10.1071/IS05031
- Philips TK, Pretorius E, Scholtz CH (2004) A phylogenetic analysis of dung beetles (Scarabaeinae: Scarabaeidae): unrolling an evolutionary history. *Invertebrate Systematics* 18: 53–88. doi:10.1071/IS03030
- Scholtz CH, Howden HF (1987) A revision of the African Canthonina. *Journal of the Entomological Society of Southern Africa* 50: 75–119.
- Sole CL, Scholtz CH (2010) Did dung beetles arise in Africa? A phylogenetic hypothesis based on five gene regions. *Molecular Phylogenetics and Evolution* 56: 631–641. doi:10.1016/j.ympev.2010.04.023

- Smith ABT (2006) A review of the family-group names for the superfamily Scarabaeoidea (Coleoptera) with corrections to nomenclature and a current classification. *Coleopterists Society Monographs* 5: 144–204.
- Vaz-de-Mello FZ (2007) *Revision taxonomica e analysis phylogenetico de la tribu Ateuchini*. PhD thesis, Mexico: Instituto de Ecología A.C, Xalapa, Veracruz.
- Vaz-de-Mello FZ (2008) Synopsis of the new subtribe Scatimina (Coleoptera: Scarabaeidae: Scarabaeinae: Ateuchini), with descriptions of twelve new genera and review of *Genieridium*, new genus. *Zootaxa* 1955: 1–75.